

Clinical Psychology

Measuring Depression and Anxiety With 4 Items? Adaptation of the PHQ-4 to Increase Its Sensitivity to Subclinical Variability

Dominique Makowski^{1,2,a}, An Shu Te³, Ana Neves¹, S.H. Annabel Chen^{3,4,5}

¹ School of Psychology, University of Sussex, Brighton, UK, ² Sussex Centre for Consciousness Science, University of Sussex, Brighton, UK, ³ School of Social Sciences, Nanyang Technological University, Singapore, ⁴ LKC Medicine, Nanyang Technological University, Singapore, ⁵ Centre for Research and Development in Learning, Nanyang Technological University, Singapore

Keywords: PHQ-4, depression, anxiety, brief questionnaire validation, ultra short scale

<https://doi.org/10.1525/collabra.143431>

Collabra: Psychology

Vol. 11, Issue 1, 2025

The PHQ-4 is an ultra-brief (4 items) screening questionnaire for depression and anxiety. In this brief report, we test the benefits of adding one additional response option (“Once or twice”, in between “Not at all” and “Several days”) to improve the scale’s sensitivity to milder alterations, and thus increase its usefulness in subclinical populations. A first pilot study (N=485) showed a prevalent usage of the new response option by participants. Its utility was confirmed in a follow-up validation study (N=836), where we that the refined version offers an improved sensitivity to subclinical variability in depression (indexed by the BDI-II) as compared to the original version. In conclusion, adding the “once or twice” response option is a low-cost way of increasing the PHQ-4’s sensitivity to subclinical variability, making it a tool of choice for general population research.

The Patient Health Questionnaire-4 (PHQ-4) is an ultra brief measurement of core signs of depression and anxiety (Kroenke et al., 2009). It consists of two items for depression (PHQ-2, Kroenke et al., 2003) and anxiety (GAD-2, Kroenke et al., 2007), each corresponding to DSM-5 diagnostic symptoms for major depressive disorder (MDD) and generalized anxiety disorder (GAD). It has been validated across many languages and populations (Christodoulaki et al., 2022; Materu et al., 2020; Mendoza et al., 2022), becoming one of the most popular screening instruments for depression and anxiety (Maurer et al., 2018).

While the scale has been validated and used in the general population and non-clinical samples (Hajek & König, 2020; Löwe et al., 2010), its initial purpose was to reliably discriminate and identify potential MDD/GAD patients. This discriminative goal materializes in the scale’s design and the existence of categorical cut-offs, which does not necessarily entail a focus on the sensitivity to milder mood alterations. In particular, the gap between the two lowest possible answers, “Not at all” and “Several days”, is quite large and possibly leaves out the possibility of more subtle occurrences. While this is not necessarily an issue in clinical and diagnostic contexts, it might lead to a sub-optimal discrimination of affective levels on the lower end of the spectrum, important for instance in the context of subclinical variability quantification.

Enabling a more precise assessment of low-severity alterations is important, as milder symptoms are significant

predictors of future clinical disorders and are associated with present functional impairments and reduced quality of life (Cuijpers & Smit, 2004; Judd et al., 1998). In addition, the growing reliance on large-scale, often online, psychological and epidemiological surveys, to monitor population-level mental health, evaluate interventions, or track responses to global stressors such as pandemics or geopolitical crises, demands tools that are both brief and sensitive to small but meaningful fluctuations.

This brief report aims at testing the possibility of enhancing - with minimal changes to the original scale - the PHQ-4 sensitivity to mild mood level inflections. In a first pilot study, we will evaluate whether the new response option is used by participants as well as its prevalence. In a follow-up validation study, we will compare the refined PHQ-4 version to the original one in terms of sensitivity to subclinical variability in depression, using the Beck Depression Inventory (BDI-II, Beck et al., 1996) and the Trait scale of the State-Trait Anxiety Inventory (STAI-5, Zsido et al., 2020) as reference measures of depression and anxiety. This refinement targets researchers and practitioners conducting large-scale or longitudinal studies, particularly in public health and psychological research, in need of brief yet sensitive instruments to track and explore mental health trends.

^a Correspondence concerning this article should be addressed to Dominique Makowski, School of Psychology, University of Sussex, Brighton BN1 9RH, UK, Email: D.Makowski@sussex.ac.uk

Study 1 (Pilot)

Method

Participants

The sample consists of 485 English-speaking participants (Mean age = 30.1 ± 10.1 [18, 73]; 50.3% females) from the general population recruited via *Prolific*, a crowd-sourcing platform recognized for providing high quality data (Peer et al., 2022). The only inclusion criterion was a fluent proficiency in English to ensure that the task instructions would be well-understood. This study was approved by the NTU Institutional Review Board (NTU IRB-2022-187). All participants provided their informed consent prior to participation and were incentivized after completing the study.

Measures

In the original PHQ-4, the instructions “Over the last 2 weeks, how often have you been bothered by the following problems?” are followed with 4 items (A1 - *Feeling nervous, anxious or on edge*; A2 - *Not being able to stop or control worrying*; D1 - *Little interest or pleasure in doing things*; D2 - *Feeling down, depressed, or hopeless*). The original answer options are “Not at all” (0), “Several days” (1), “More than half the days” (2), “Nearly every day” (3). The total score is computed by summing the responses of each facet resulting in a 0-6 score for depression and anxiety.

For the refined version, we added a “Once or twice” option between “Not at all” and “Several days” in order to better capture potential mild mood inflections (see Dobson & Mothersill, 1979 for the choice of the label).

Procedure

Participants were administered the refined PHQ-4 online as part of another study, which contained additional questionnaires and tasks not relevant for the current analysis. The PHQ-4 was presented in a randomized order with other questionnaires. The data is available in open-access at <https://github.com/RealityBending/IlusionGameReliability>.

Results

The analysis was carried out using *R 4.4* (R Core Team, 2023), the *tidyverse* (Wickham et al., 2019), and the *easystats* collection of packages (Lüdtke et al., 2019, 2020, 2021; Patil et al., 2022). All reproducible scripts and complementary analyses are available open-access at <https://github.com/DominiqueMakowski/PHQ4R>

The consistency of the anxiety (*Cronbach's* $\alpha = 0.903$) and depression (*Cronbach's* $\alpha = 0.841$) subscales is excellent (responses were treated as ordinal). The proportion of response types stratified by item (see [Figure 1](#)) shows that the new “Once or twice” option was the most prevalent response for all items (on average selected in 29.12% of cases).

Discussion

The fact that the new “Once or twice” response option was the most prevalent response shows that it captures a meaningful (with respect to what participants’ subjective experience) degree of mood expression. The fact that this response option appears as a popular choice begs the question of its usefulness in capturing more fine-grained variations of the underlying dimensions (depression and anxiety) as measured by independent tools, which was the goal of the follow-up validation study.

Study 2 (Validation)

Method

Participants

The initial sample consisted of 1053 participants, recruited (181 were recruited on *Prolific*, 772 students from the University of Sussex via *SONA*, and the rest through convenience sampling as part of dissertation students’ data collection). We used attention checks as the primary target for participant exclusion. We excluded 194 participants (18.42%) for failing at least one attention check, and 23 (2.18%) that were outliers ($|z_{robust}| > 2.58$) on measures significantly related to the probability of failing attention checks (namely, the standard deviation of all the items of the Interoceptive Accuracy Scale, as well as the the multivariate distance obtained with the OPTICS algorithm, see Thériault et al., 2024). The experiment duration was not related to the probability of failing attention checks and was thus not used as an exclusion criterion.


The final sample included 836 participants (Mean age = 25.1 ± 11.3 [18, 76]; 73.8% women). This study was approved by the University of Sussex’ Ethics Committee (ER/ASF25/4).

In this sample, 51 participants (6.10%) were coded as having Depression, as indexed by the self-reported presence of MDD together with the use of a treatment (antidepressant, anxiolytic and/or therapy), and 87 participants (10.41%) were coded as having Anxiety, as indexed by the self-reported presence of GAD or Panic Disorder, also together with the use of a treatment.

Measures

Participants were randomly assigned to complete either the original or refined version of the PHQ-4, which included one additional response option (“Once or twice”). To preserve comparability with the original PHQ-4 scoring system and avoid altering the scale’s total score range or established cut-off thresholds, we assigned the new response option a value of 0.5, placing it midway between “Not at all” (0) and “Several days” (1). While this scoring assumes equal spacing between response options - a common but imperfect convention in ordinal scales - it offers a pragmatic compromise between conceptual fidelity and applied utility.

A. Proportion of answers per item



B. Joint prevalence of responses




Figure 1. Prevalence of response options to the refined version of the PHQ-4. A) Proportion of answers of each type to the four items. B) Prevalence of all pairs of answers.

Beck's Depression Inventory (BDI-II, Beck et al., 1996) was used as a reference measure of depressive symptoms. It includes 21 items, each addressing a specific depression symptom and offering four response options scored from 0 to 3. Participants are instructed to select the option that best describes how they have felt over the past two weeks. The total score is calculated by summing the scores for all 21 items, with higher scores indicating greater severity of depressive symptoms.

The short version of the trait subscale of the State-Trait Anxiety Inventory (STAI-5, Zsido et al., 2020) was used as a reference measure of anxiety. This abridged version of the STAI (Spielberger, 1970) includes 5 items rated on a 4-point Likert scale. Changes were made in the instructions from asking "how participants feel right now" to "over the past 2 weeks" to keep it consistent with the instructions of the PHQ-4 and BDI-II. A general score of anxiety was computed by averaging all the items.

Participants were also asked to complete two questionnaires of interoception, namely the Interoceptive Accuracy Scale (IAS, 21 items rated on analog scales, Murphy et al., 2020) and the Multidimensional Assessment of Interoceptive Awareness (MAIA-2 - 37 items, Mehling et al., 2018).

After demographic questions, participants were asked to report the current presence of psychiatric issues (from a list), as well as the usage of treatment (antidepressants, mood stabilizers, anxiolytics, therapy). We indexed the presence of a depression when participants reported suffering from either Major Depressive Disorder (MDD) or Dysthymia, as well as undergoing a medical treatment. Similarly, we indexed the presence of an anxiety disorder when participants reported suffering from either Generalized Anxiety Disorder (GAD) or Panic Disorder, as well as undergoing a medical treatment.

Procedure

The original or refined version of the PHQ-4 was followed by the BDI-II, STAI-5, IAS, and MAIA-2, presented in random order. The IAS (Murphy et al., 2020) and the MAIA-2 (Mehling et al., 2018) were included as part of another focused on interoception, and were only used in this study as part of data quality control checks.

Results

As all the scripts, analysis details and results tables are available open-access at <https://github.com/Dominique-Makowski/PHQ4R>, we will focus on reporting the main results.

Bayesian *t*-tests (using *BayesFactor*'s *ttestBF()* function with default priors, Morey & Rouder, 2024) comparing the two groups ("original version", *N* = 396 vs. "refined version", *N* = 440) revealed evidence against the existence of a difference for the BDI-II total score ($BF_{10} = 0.20$) and the STAI-5 general score ($BF_{10} = 0.08$), suggesting their baseline comparability in mood level. Moreover, the two groups did also not differ in their PHQ-4 depression ($BF_{10} = 0.23$) and anxiety ($BF_{10} = 0.09$) scores, despite the differences in scoring introduced by the refined version, suggesting their comparability.

PHQ-4 Depression vs. BDI-II

A linear model testing the interaction effect Δ of the refined condition on the intercept (representing by how much the value of the outcome when the PHQ score is 0 changes for the refined version compared the original) and slope (its increase or decrease by the refined version compared to the original) of the relationship between the PHQ-4 depression score and the BDI-II total score was fitted. The model predicting the BDI-II total score with the PHQ-4 depression score showed no interaction related to the PHQ-4 version ($\Delta Intercept_{\text{refined}} = -0.13$, 95% *CI* [-1.73, 1.47], $t(832) = -0.16$, $p = 0.871$; $\Delta \beta_{\text{refined}} = -0.05$, 95% *CI* [-0.70, 0.60], $t(832) = -0.15$, $p = 0.883$), suggesting no differences in the relationship pattern between the two versions (see [Figure 2](#)).

Moreover, Bayesian *t*-tests comparing the BDI-II scores between the refined and the original version at each integer score (0, 1, 2, 3) yielded no evidence in favour of a significant difference ($BFs < 3$). In other words, having the same score on the refined version as on the original version was related to the same outcome on the BDI-II (further evidencing the comparability between the 2 versions).

However, the low in-between scores from the refined version are overall capturing significantly different levels of depression compared to the adjacent scores. Scoring 0.5 was associated with a higher BDI-II score than scoring 0 ($BF > 30$), and lower scores than scoring 1 ($BF > 30$). Similarly, scoring 1.5 was associated with a higher BDI-II score than scoring 1 ($BF > 30$), but not lower scores than scoring 2 ($BF = 0.234$).

PHQ-4 Anxiety vs. STAI-5

The linear regression predicting the STAI-5 general score with the PHQ-4 anxiety score showed no interaction related to the PHQ-4 version ($\Delta Intercept_{\text{refined}} = -0.02$, 95% *CI* [-0.15, 0.11], $t(832) = -0.32$, $p = 0.750$; $\Delta \beta_{\text{refined}} = 0.01$, 95% *CI* [-0.03, 0.05], $t(832) = 0.56$, $p = 0.576$), suggesting no differences in the relationship pattern between the two versions.

Moreover, Bayesian *t*-tests comparing the STAI-5 scores between the refined and the original version at each integer score yielded no evidence in favour of a significant difference. In other words, having the same score on the refined version as on the original version was related to the same outcome on the STAI-5.

However, comparing in-between scores with adjacent scores yielded mixed results. Scoring 0.5 on the PHQ-4 anxiety was not significantly associated with a different level of STAI-5 compared to scoring 0 ($BF = 1.83$), but was with scores of 1 ($BF > 30$). Similarly, there was no evidence that scoring 1.5 was different from scoring 1 ($BF = 0.605$), but strong evidence that it was different from scoring 2 ($BF > 30$).

Correlation Differences

While the relationship pattern (i.e., the slope of the linear relationship) was not affected by the PHQ-4 version, we focused next on testing the difference in the strength (i.e., the precision) of the relationship, in particular at the lower end of the spectrum (i.e., for sub-clinical threshold scores of the BDI-II and STAI-5). We bootstrapped (2000 iterations) the difference in correlation between the refined and the original version for each of the two reference measures, separately for the BDI-II subsamples (minimal to mild ≤ 18 ; moderate to severe > 18) and the STAI-5 subsamples (minimal to mild < 2 ; moderate to severe ≥ 2).

The results suggested that in the subclinical range of the BDI-II, the correlation between its score and the PHQ-4 Depression score was marginally higher (although not significantly, $p_{\text{one-sided}} = 0.164$) for the refined version compared to the original one. No correlation differences were observed in the moderate to severe range of the BDI-II.




Figure 2. A) PHQ-4 depression and anxiety scores against their respective reference measures, the BDI-22 and the STAI-5. Bayes factors in grey tell if there is a difference, for the same PHQ-4 score, between the original and the refined version (BFs < 1 suggest no difference and thus evidence for a comparability of the refined version with respect to the original scale. Bayes factors in yellow represent how new in-between scores (0.5, 1.5, 2.5, ...) available with refined version differ from the adjacent scores (BFs > 3 suggest that half a point of difference on the refined PHQ-4 relates to a significant difference on the reference measure). BF < 1/3°, BF > 3°, BF > 10**, BF > 30***. B) Bootstrapped distributions of the difference of correlation between the revised PHQ-4 scores and the original one for sub-clinical threshold scores of depression and anxiety. Positive differences suggest that the correlation between the reference measure and the refined PHQ-4 score was stronger compared to the original version. C) Predictive power of the PHQ-4 scores on the presence of a depression or anxiety disorder. The upper plots show the relationship modelled by a logistic regression, while the above plots represent the ROC curves (in which a line further away from the diagonal represents a higher combination of sensitivity and specificity).

For the STAI-5, there was no difference in the correlation between the refined and the original version in the sub-clinical range of the STAI-5. Surprisingly, we observed a stronger correlation between the refined PHQ-4 Anxiety score and the STAI-5 in the moderate to severe range compared to the original version ($p_{one-sided} = 0.017$).

Predictive Power

Finally, we tested the predictive power of the PHQ-4 depression and anxiety scores on the presence of a depression or anxiety disorder, respectively. We modeled the relationship with a logistic regression. While the PHQ-4 was overall a strong predictor of the outcome, there was no significant difference between the two PHQ-4 versions.

However, the ROC curves for the refined and the original version of the PHQ-4, suggested that the refined version had a better sensitivity / specificity trade-off (AUC = 78.36%) compared to the original version (AUC=75%), in particular on the lower end of the spectrum. The difference was negligible for anxiety.

Discussion

These results suggest that the new “Once or twice” response option to the PHQ-4 does help capturing more fine-grained variations of depressive symptoms, particularly in the subclinical range. Importantly, adding this new response option with the scoring of 0.5 does not disrupt the quality of the scale, which scores remain comparable to that of the original version.

The results for the anxiety subscale appear more mixed, with less evident benefits. However, this might have been partly caused by our design decision regarding the questionnaire used for the reference measure of anxiety. Indeed, we used the abridged version of the STAI, which only included 5 items, arguably limiting the sensitivity of the anxiety measure in the first place.

One of the potential limitations of our study includes the choice of the measures used as external reference proxies of depression and anxiety, namely the BDI-II and the STAI-5. While both scales are widely used, it is important to underline that they are not perfect and pure measures of depression and anxiety. For instance, there is ongoing debate about the STAI’s discriminant validity (which might extend to its short form used in the present study). A recent meta-analysis reported that trait anxiety scores were more strongly associated with depressive than anxiety disorders (Knowles & Olatunji, 2020), suggesting that the scale may rather capture general negative affect than specifically anxiety. Regarding the BDI-II, existing evidence that suggests its relative lack of sensitivity for low-severity depression (Olino et al., 2012) - comparable to similar instruments of its size (e.g., CES-D), and an improvement over the BDI-I (Wahl et al., 2014) - put into question its choice for our goal of showing increased sub-clinical sensitivity. While we believe that evidence in favour of differences found for a possibly insensitive measure does not critically compromise our conclusions (as opposed to finding no differences, which could be explained by the low sensitivity), future

studies should verify these findings with alternative measures of anxiety and depression and clinically assessed populations.

Indeed, although we used a stricter criterion for classifying participants as having a depression or an anxiety disorder by restricting it to participants also reporting undergoing a medical treatment, it was still based on self-reported data. Studies in controlled clinical settings are needed to confirm the potential benefits of the refined PHQ-4 in mood disorders detection accuracy.

General Discussion

The objective of this study was to test the introduction of a “Once or twice” response option to the PHQ-4 to enhance its sensitivity to milder mood fluctuations. In the first pilot study, we showed that the new response option was used prevalently, suggesting relevance and increased precision for participants to express their mood difficulties. The follow-up validation study showed that the refined version of the PHQ-4 was able to better differentiate lower levels of depression compared to the original version, while remaining comparable. Although the benefits of this refinement appear to be fairly minor, and particularly marked for the depression score compared to anxiety, this low-cost improvement appear useful to implement when measuring depression and anxiety using the PHQ-4 ultra-short screening questionnaire.

While adding granularity to the response format appears useful for the PHQ-4, it is possible that similar benefits could be found with other scales and measures. While the use of a limited, highly discriminating set of response options is understandable for specific applications (e.g., clinical diagnosis), we recommend future studies to investigate response format and its potential improvement for other scales used in online surveys and general population research.

Author Contributions

Author roles were classified using the Contributor Role Taxonomy (CRediT; <https://credit.niso.org/>) as follows: *Dominique Makowski*: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, software, supervision, validation, visualization, and writing – original draft. *An Shu Te*: project administration, resources, and writing – original draft. *Ana Neves*: data curation, formal analysis, writing – original draft, and writing – review & editing. *S.H. Annabel Chen*: project administration, supervision, and writing – review & editing

Competing Interests

The authors declare no conflict of interest.

Data Accessibility

Data and reproducible analyses are available at <https://github.com/DominiqueMakowski/PHQ4R>.

Editors: Daniel P. Moriarity (Associate Editor)

Submitted: January 23, 2025 PDT. Accepted: August 06, 2025 PDT. Published: August 22, 2025 PDT.

Acknowledgements

We would like to thank the dissertation students from the University of Sussex for their help in data collection.



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

References

- Beck, A. T., Steer, R. A., ... Brown, G. K. (1996). *Beck depression inventory*. <https://doi.org/10.1037/t00742-000>
- Christodoulaki, A., Baralou, V., Konstantakopoulos, G., & Touloumi, G. (2022). Validation of the patient health questionnaire-4 (PHQ-4) to screen for depression and anxiety in the greek general population. *Journal of Psychosomatic Research*, 160, 110970. <https://doi.org/10.1016/j.jpsychores.2022.110970>
- Cuijpers, P., & Smit, F. (2004). Subthreshold depression as a risk indicator for major depressive disorder: A systematic review of prospective studies. *Acta Psychiatrica Scandinavica*, 109(5), 325–331. <https://doi.org/10.1111/j.1600-0447.2004.00301.x>
- Dobson, K. S., & Mothersill, K. J. (1979). Equidistant categorical labels for construction of likert-type scales. *Perceptual and Motor Skills*, 49(2), 575–580. <https://doi.org/10.2466/pms.1979.49.2.575>
- Hajek, A., & König, H.-H. (2020). Prevalence and correlates of individuals screening positive for depression and anxiety on the phq-4 in the german general population: Findings from the nationally representative german socio-economic panel (GSOEP). *International Journal of Environmental Research and Public Health*, 17(21), 7865. <https://doi.org/10.3390/ijerph17217865>
- Judd, L. L., Akiskal, H. S., Maser, J. D., Zeller, P. J., Endicott, J., Coryell, W., Paulus, M. P., Kunovac, J. L., Leon, A. C., ... Mueller, T. I. (1998). Major depressive disorder: A prospective study of residual subthreshold depressive symptoms as predictor of rapid relapse. *Journal of Affective Disorders*, 50(2–3), 97–108. [https://doi.org/10.1016/S0165-0327\(98\)00138-4](https://doi.org/10.1016/S0165-0327(98)00138-4)
- Knowles, K. A., & Olatunji, B. O. (2020). Specificity of trait anxiety in anxiety and depression: Meta-analysis of the state-trait anxiety inventory. *Clinical Psychology Review*, 82, 101928. <https://doi.org/10.1016/j.cpr.2020.101928>
- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2003). The patient health questionnaire-2: Validity of a two-item depression screener. *Medical Care*, 1284–1292. <https://doi.org/10.1097/01.MLR.0000093487.78664.3C>
- Kroenke, K., Spitzer, R. L., Williams, J. B., & Löwe, B. (2009). An ultra-brief screening scale for anxiety and depression: The PHQ-4. *Psychosomatics*, 50(6), 613–621.
- Kroenke, K., Spitzer, R. L., Williams, J. B., Monahan, P. O., & Löwe, B. (2007). Anxiety disorders in primary care: Prevalence, impairment, comorbidity, and detection. *Annals of Internal Medicine*, 146(5), 317–325. <https://doi.org/10.7326/0003-4819-146-5-200703060-00004>
- Löwe, B., Wahl, I., Rose, M., Spitzer, C., Glaesmer, H., Wingenfeld, K., Schneider, A., & Brähler, E. (2010). A 4-item measure of depression and anxiety: Validation and standardization of the patient health questionnaire-4 (PHQ-4) in the general population. *Journal of Affective Disorders*, 122(1–2), 86–95. <https://doi.org/10.1016/j.jad.2009.06.019>
- Lüdtke, D., Ben-Shachar, M. S., Patil, I., & Makowski, D. (2020). Extracting, computing and exploring the parameters of statistical models using r. *Journal of Open Source Software*, 5(53), 2445. <https://doi.org/10.21105/joss.02445>
- Lüdtke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., & Makowski, D. (2021). Performance: An r package for assessment, comparison and testing of statistical models. *Journal of Open Source Software*, 6(60). <https://doi.org/10.21105/joss.03139>
- Lüdtke, D., Waggoner, P. D., & Makowski, D. (2019). Insight: A unified interface to access information from model objects in r. *Journal of Open Source Software*, 4(38), 1412. <https://doi.org/10.21105/joss.01412>
- Materu, J., Kuringe, E., Nyato, D., Galishi, A., Mwanamsangu, A., Katebalila, M., Shao, A., Chagalucha, J., Nnko, S., & Wambura, M. (2020). The psychometric properties of PHQ-4 anxiety and depression screening scale among out of school adolescent girls and young women in tanzania: A cross-sectional study. *BMC Psychiatry*, 20(1), 1–8. <https://doi.org/10.1186/s12888-020-02735-5>
- Maurer, D. M., Raymond, T. J., & Davis, B. N. (2018). Depression: Screening and diagnosis. *American Family Physician*, 98(8), 508–515.
- Mehling, W. E., Acree, M., Stewart, A., Silas, J., & Jones, A. (2018). The multidimensional assessment of interoceptive awareness, version 2 (MAIA-2). *PloS One*, 13(12), e0208034. <https://doi.org/10.1371/journal.pone.0208034>
- Mendoza, N. B., Frondoza, C. E., Dizon, J. I. W. T., & Buenconsejo, J. U. (2022). The factor structure and measurement invariance of the PHQ-4 and the prevalence of depression and anxiety in a southeast asian context amid the COVID-19 pandemic. *Current Psychology*, 1–10.
- Morey, R. D., & Rouder, J. N. (2024). *BayesFactor: Computation of bayes factors for common designs*. <https://CRAN.R-project.org/package=BayesFactor>
- Murphy, J., Brewer, R., Plans, D., Khalsa, S. S., Catmur, C., & Bird, G. (2020). Testing the independence of self-reported interoceptive accuracy and attention. *Quarterly Journal of Experimental Psychology*, 73(1), 115–133. <https://doi.org/10.1177/1747021819879826>
- Olino, T. M., Yu, L., Klein, D. N., Rohde, P., Seeley, J. R., Pilkonis, P. A., & Lewinsohn, P. M. (2012). Measuring depression using item response theory: An examination of three measures of depressive symptomatology. *International Journal of Methods in Psychiatric Research*, 21(1), 76–85. <https://doi.org/10.1002/mp.1348>

- Patil, I., Makowski, D., Ben-Shachar, M. S., Wiernik, B. M., Bacher, E., & Lüdtke, D. (2022). Datawizard: An r package for easy data preparation and statistical transformations. *Journal of Open Source Software*, 7(78), 4684. <https://doi.org/10.21105/joss.04684>
- Peer, E., Rothschild, D., Gordon, A., Evernden, Z., & Damer, E. (2022). Data quality of platforms and panels for online behavioral research. *Behavior Research Methods*, 54(4), 1643–1662. <https://doi.org/10.3758/s13428-021-01694-3>
- R Core Team. (2023). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Spielberger, C. D. (1970). *Manual for the state-trait anxiety inventory (self-evaluation questionnaire)*.
- Thériault, R., Ben-Shachar, M. S., Patil, I., Lüdtke, D., Wiernik, B. M., & Makowski, D. (2024). Check your outliers! An introduction to identifying statistical outliers in r with easystats. *Behavior Research Methods*, 56(4), 4162–4172. <https://doi.org/10.3758/s13428-024-02356-w>
- Wahl, I., Löwe, B., Bjorner, J. B., Fischer, F., Langs, G., Voderholzer, U., Aita, S. A., Bergemann, N., Brähler, E., & Rose, M. (2014). Standardization of depression measurement: A common metric was developed for 11 self-report depression measures. *Journal of Clinical Epidemiology*, 67(1), 73–86. <https://doi.org/10.1016/j.jclinepi.2013.04.019>
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Golemund, G., Hayes, A., Henry, L., ... Hester, J. (2019). Welcome to the tidyverse. *Journal of Open Source Software*, 4(43), 1686. <https://doi.org/10.21105/joss.01686>
- Zsido, A. N., Teleki, S. A., Csokasi, K., Rozsa, S., & Bandi, S. A. (2020). Development of the short version of the spielberger state–trait anxiety inventory. *Psychiatry Research*, 291, 113223. <https://doi.org/10.1016/j.psychres.2020.113223>

Supplementary Materials

Peer Review Communication

Download: https://collabra.scholasticahq.com/article/143431-measuring-depression-and-anxiety-with-4-items-adaptation-of-the-phq-4-to-increase-its-sensitivity-to-subclinical-variability/attachment/298725.docx?auth_token=AAD3nucKOIW9O7w77GYr
